

Quantum Communication Protocol Employing Weak Measurements

Alonso Botero^{1,*} and Benni Reznik²

¹ Center for Particle Physics, University of Texas, Austin, Texas 78712

² School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel

We propose a communication protocol exploiting correlations between two events with a definite time-ordering: a) the outcome of a *weak measurement* on a spin, and b) the outcome of a subsequent ordinary measurement on the spin. In our protocol, Alice, first generates a “code” by performing weak measurements on a sample of N spins. The sample is sent to Bob, who later performs a post-selection by measuring the spin along either of two certain directions. The results of the post-selection define the “key”, which he then broadcasts publicly. Using both her previously generated code and this key, Alice is able to infer the *direction* chosen by Bob in the post-selection. Alternatively, if Alice broadcasts publicly her code, Bob is able to infer from the code and the key the direction chosen by Alice for her weak measurement. Two possible experimental realizations of the protocols are briefly mentioned.

Weak measurements [1–3] are a special class of quantum measurements explored in recent years by Aharonov, Vaidman, and others. In one such measurement of any given observable A , the disturbance caused to the system is minimized at the expense of precision in a single trial. Nevertheless, after a large number trials one can determine statistical averages such as the expectation value of A . The distinctive feature of weak measurements has to do with the observed averages when the measured system is *post-selected*. Such averages, the so-called *weak values*, may lie outside the bounds of the spectrum of A [2]. Moreover, since they depend on the chosen pre- and post-selected ensemble, weak values carry non-trivial information about the *choice* of measurement used for post-selection. In accordance with causality, these unusual regularities must therefore be *a priori* undetectable, i.e., “hidden in the noise” [3]. Hence, they can only be extracted *a posteriori*, from the correlations between readings of the measurement and the result of the post-selection.

In the two protocols suggested in this Letter, such correlations are used for signaling. In the first protocol, the receiver, Alice, starts by generating a *code* corresponding to the outcomes of a series of weak measurements on a large sample of spins. She then hands them to Bob, who is about to depart on a long voyage, together with an important question that he can only answer at some later time. When he is ready to respond, Bob performs an ordinary measurement (post-selection) on these spins. His response corresponds to the choice of spin component he then measures; for instance, a measurement of σ_y to signal a “yes” (“I will marry you”), or a measurement of σ_z to signal a “no”. The random sequence of results obtained in either of these measurements plays the role

of a *key*, which he then broadcasts publicly. This is just enough information for Alice to bin her previous readings and extract the message from the weak values. The random key is useless to the ever-jealous Eve; it can be shown that an optimal sample size exists which is still too small for Eve to infer Bob’s choice from the statistics of the key.

The security of this protocol is however compromised if the spins fall in the hands of the eavesdropper. This drawback is overcome in the second protocol, in which instead it is Alice who *sends* the message; for this, she performs on the N spins either of two possible weak measurements and generates a code corresponding to her outcomes. She then sends the spins *and* the code to Bob. To decode the message Bob post-selects the spins by performing a sequence of strong measurements. As in the first case, the resulting key allows him to bin the code and deduce from the weak values Alice’s choice of measurement. In this case, Eve cannot intercept the message from the spins without Bob being aware of her actions.

We begin by analyzing the weak measurement scheme. Suppose that Alice prepares a sample of N spin-1/2 particles, in the eigenstate $|x+\rangle$ of σ_x , and she wishes to perform a weak measurement of the spin observable

$$A \equiv \frac{1}{\sqrt{2}}(\sigma_x + \sigma_y), \quad (1)$$

with eigenstates $|a\pm\rangle$. We let p be the pointer variable of the measuring device and $|\phi_i\rangle$ its initial state, with a Gaussian wave function $\phi_i(p) = (2\pi\Delta p^2)^{-1/4}e^{-p^2/4\Delta p^2}$ of uncertainty Δp . The weak measurement may then be described by the usual transformation taking an initial product state $|\Psi_i\rangle \equiv |x+\rangle|\phi_i\rangle$ of the spin and the device, to a final entangled state:

$$|\Psi_i\rangle \rightarrow |\Psi_f\rangle = c_+|a+\rangle|\phi+\rangle + c_-|a-\rangle|\phi-\rangle, \quad (2)$$

where $c_{\pm} = \langle a \pm | x + \rangle$, and $|\phi\pm\rangle$ is $|\phi_i\rangle$ shifted in p by ± 1 . In contrast however to an ordinary measurement, Δp is here assumed to be so large that the shifted states $|\phi\pm\rangle$ overlap considerably. Nevertheless, the expectation value of p is still shifted by the usual expectation value $\langle A \rangle$:

$$\langle \Psi_f | p | \Psi_f \rangle = \langle x + | A | x + \rangle = \frac{1}{\sqrt{2}}. \quad (3)$$

The idea of a weak measurement is thus to extract such systematic shifts of the means from a large sample of identically prepared spins, ensuring at the same time a

minimal disturbance of any individual spin. This disturbance is naturally related to the overlap between $|\phi+\rangle$ and $|\phi-\rangle$.

A convenient measure of disturbance D is given in terms of the Fidelity $\mathcal{F}$ [4] as $D \equiv 1 - \mathcal{F}$, where $\mathcal{F}$ is defined as the probability of obtaining back the initial state after the weak measurement, $\mathcal{F} = ||\langle x + |\Psi_f\rangle||^2$; hence, D may be interpreted as the probability of “flipping” the initial direction of the spin. As one verifies, D involves the overlap factor $\langle \phi + | \phi - \rangle = e^{-1/2\Delta p^2}$:

$$D = \frac{\Delta A^2}{2}(1 - e^{-1/2\Delta p^2}) \simeq \frac{1}{8\Delta p^2}. \quad (4)$$

We assume in the approximation that D is small and take $\Delta A^2 = \langle A^2 \rangle - \langle A \rangle^2 = 1/2$.

Together then with the weakness condition $D \ll 1$, the resulting distribution for the pointer variable must be sufficiently broad that the spectrum of A cannot be resolved. The distribution therefore takes essentially a Gaussian form, centered at the expectation value $\langle A \rangle$, with a large uncertainty $\sqrt{\Delta p^2 + \Delta A^2} \simeq \Delta p$ of order $D^{-1/2}$. Since in N identical and independent trials the expected sample mean is $\bar{p} = \langle A \rangle \pm \Delta p/\sqrt{N}$, Alice can determine $\langle A \rangle$ if her sample is large enough.

But now suppose that after completing her measurements Alice gives the spins to Bob, who later performs a second, now ordinary measurement on each spin, along some arbitrary direction $\hat{b}$ with possible outcomes $|b\pm\rangle$. Each outcome in this *post-selection* is correlated with a particular response of the measuring device. To keep track of these correlations it is convenient to re-express the combined state of the system and apparatus in terms of the basis $|b\pm\rangle$

$$|\Psi_f\rangle = |b+\rangle|\phi(b+)\rangle + |b-\rangle|\phi(b-)\rangle, \quad (5)$$

where the conditional response of the apparatus is described by the unnormalized states $|\phi(b\pm)\rangle = \langle b\pm | \Psi_f \rangle$. For a given result $|b\rangle$,

$$|\phi(b)\rangle = C \left[\frac{1 + A_w(b)}{2} |\phi+\rangle + \frac{1 - A_w(b)}{2} |\phi-\rangle \right], \quad (6)$$

here $C = \langle b|x+\rangle$ and A_w is the *weak value* of A , defined as

$$A_w(b) \equiv \frac{\langle b|A|x+\rangle}{\langle b|x+\rangle}. \quad (7)$$

The weak value $A_w(b)$ is the defining property of weak measurements: as before, when D is small the pointer distribution is essentially Gaussian of uncertainty $\simeq \Delta p$; however, as one can easily show, the location of the mean $\langle p \rangle = \frac{\langle \phi(b)|p|\phi(b) \rangle}{\langle \phi(b)|\phi(b) \rangle}$ is now given by

$$\langle p \rangle = \frac{\text{Re}A_w(b)}{1 + \frac{\delta P(b)}{P(b)}} = \text{Re}A_w + O(D) \quad (8)$$

where, in terms of D ,

$$\frac{\delta P(b)}{P(b)} = -\frac{1 - |A_w(b)|^2}{\Delta A^2} D. \quad (9)$$

The correction $\frac{\delta P(b)}{P(b)}$ is the relative deviation of the probability $P'(b) = \langle \phi(b)|\phi(b) \rangle$ of obtaining $|b\rangle$ in the presence of the weak measurement, as measured from the unperturbed probability $P(b) = |\langle b|\psi \rangle|^2$. We see therefore that the average conditional reading becomes the real part of the weak value $A_w(b)$ as we approach the ideally weak situation in which the transition probability is left unaltered by the measurement.

To summarize then: if Alice starts with a large sample of N identical spins, she obtains a broad distribution of pointer readings with a sample mean $\langle A \rangle \pm \Delta p/\sqrt{N}$; however, were she to know—*for every single spin*—which of the two possible outcomes $|b+\rangle$ and $|b-\rangle$ was obtained in Bob’s measurement, then she could divide her readings into two categories, corresponding to two post-selected sub-samples of size $N_{\pm} \simeq NP(b_{\pm})$. Her original distribution would then break up as a mixture of two conditional distributions, with sample means $A_w(b+)$ and $A_w(b-)$ within errors of $\Delta p/\sqrt{N_{\pm}}$. The break-up is captured by a simple sum rule [5] in the limit $D \rightarrow 0$

$$\langle A \rangle = P(b) \text{Re}A_w(b) + P(b-) \text{Re}A_w(b-), \quad (10)$$

which serves as the basis for our protocol.

For simplicity let us first consider a complete cycle at the end of which Alice receives from Bob a single-bit “yes” or “no” message. When Alice prepares her spin sample in the eigenstate $|x+\rangle$, she labels each spin as $i = 1, \dots, N$. From her N weak measurements of $\frac{1}{\sqrt{2}}(\sigma_x + \sigma_y)$, she then generates the code by recording a string of real numbers corresponding to the pointer readings $\{p_1, p_2, \dots, p_N\}$; the sample mean $\bar{p} = \frac{1}{N} \sum_i p_i$ should yield the expectation value $\langle A \rangle \simeq 1/\sqrt{2}$. Next the spins are given to Bob, carefully keeping track of the ordering. Now it’s his turn to send the message. If Bob decides to send a “yes”, he measures σ_y on every single spin; otherwise he measures σ_z to signal a “no”. In either case, from the resulting sequence of outcomes he generates the key: an ordered list of N bits $\{k_1, k_2, \dots, k_n\}$ where, say, $k_i = 1$ and 0 respectively correspond to the outcomes “up” and “down” for the i -th spin. It is this key which Bob sends back to Alice using an insecure channel. As the key is effectively random by virtue of the fact that $\hat{y}$ and $\hat{z}$ lie on the equal-probability plane, the uncorrelated sequence of “1” and “0”’s will be useless to the eavesdropper.

On the other hand, Alice, upon receiving the key, may go back to her code and separate each reading p_i into either of two bins, depending on whether $k_i = 1$ or 0. Finally, she computes the mean values of p in each bins

$$\bar{p}_1 = \frac{\sum_i p_i k_i}{\sum_i k_i}, \quad \bar{p}_0 = \frac{\sum_i p_i (1 - k_i)}{\sum_i (1 - k_i)}. \quad (11)$$

She takes these values as estimates of the “true” means within errors of order $\Delta p/\sqrt{N}/2$. Now, if indeed Bob sent a “yes”, then she should see that $\bar{p}_0 \simeq A_w(y-) = 0$, and an “eccentric” weak value $\bar{p}_1 \simeq A_w(y+) = \sqrt{2}$. Instead, if Bob sent a “no”, then she will find no significant deviation in either of the two means $\bar{p}_1$ or $\bar{p}_0$; for this case, the real part of $A_w(z\pm)$ coincides with the sample mean of $\langle A \rangle = 1/\sqrt{2}$. Thus, by distinguishing between a non-trivial and a trivial break-up of the sample mean, Alice can decode Bob’s single-bit message [6].




FIG. 1. Schematic of a cycle from which Alice receives a “yes” response from Bob.

One may object that due to the small disturbance of the weak measurements, the average numbers of “1”’s and “0”’s in the key are no longer identical for the two final measurements of σ_y and σ_z . Indeed, by eq. (9), there is no change in the probabilities for the measurement along z , but for the measurement along y the relative change in the probabilities is $\delta P(y\pm)/P(y\pm) = \pm 2D$. The difference may then be used by Eve to distinguish between the two messages if the string is sufficiently long. This leads us to compare the minimal sample sizes N_E and N_A , respectively, required by Eve and Alice to extract the message from their available data. Alice needs to distinguish between the statistics with means $\sqrt{2}$ and 0. Hence she requires that $\sqrt{2} \gtrsim \Delta p/\sqrt{N_A}/2$, which by eq. (4) yields $N_A \gtrsim 1/D$. On the other hand, Eve must distinguish between the relative frequencies of “1”’s of $1/2$ for “no” and $1/2 \pm D$ for “yes”, with an uncertainty in the frequencies $\Delta f \simeq 1/2\sqrt{N}$; for this she needs $N_E \gtrsim 1/D^2$ trials. The ratio between the two optimal sample sizes therefore scales like $\frac{N_E}{N_A} \sim \frac{1}{D}$ and consequently, for a sufficiently small D , an optimal value N may be chosen such that $N_E \gg N \geq N_A$ ensuring that the key is safe.

While this protocol is particularly interesting from the conceptual point of view as we further discuss below, it would fail to meet usual cryptography security requirements had Alice sent the spins to Bob using a public

channel. In such case, Eve could have intercepted the message by performing weak measurements and using the public key sent by Bob [7]. The security drawback is avoided in the second protocol, where by a similar set of measurements, Alice sends a message to Bob.

As before, Alice starts with N spins in the $|x+\rangle$ state. To send a “yes” she next performs, on each spin separately, a weak measurement of $(\sigma_x + \sigma_y)/\sqrt{2}$. To send a “no” she measures weakly $(\sigma_x - \sigma_y)/\sqrt{2}$. In either case she records the results as a code of N real numbers, and sends to Bob publicly *both* the code and the spins. To decode the message Bob decides in a *random* manner to perform on each spin an ordinary measurement, of either σ_x or σ_y . The σ_x measurement is needed as a “security check”, while the results of the σ_y measurement will be used to decode the message. If the spins were not disturbed after Alice’s measurement, in $\sim N/2$ cases that Bob measures σ_x he gets $\sigma_x = 1$ and only $\sim ND \ll N/4$ times $\sigma_x = -1$. On the other hand, he finds $\sigma_y = \pm 1$ with nearly equal probability. The latter are used to bin the code, into two subsamples each with $\sim N/4$ spins, corresponding to the post-selection of $\sigma_y = 1$ and $\sigma_y = -1$. Finally he computes the mean value of each subsample. If Alice had measured $(\sigma_x + \sigma_y)/\sqrt{2}$, Bob will find the weak values $\sqrt{2}$ and 0, respectively. If instead she had measured $(\sigma_x - \sigma_y)/\sqrt{2}$, he will find the same means but in reverse order.

To discuss the question of security, notice that the weak measurements rotate the spin to two slightly different spin states. For a typical reading $p = 1/\sqrt{2} \pm \Delta p$, the overlap between the two rotated states is of the order of $1 - D$, as they lie at angle of $\sim \sqrt{D}$ from $x+$. Now, since Bob requires $N \sim 1/D$ to decode the message, the overlap between the two “yes/no” *collective* N -spin states becomes nearly orthogonal. The crucial point is that now there are two distinct ways to distinguish between these states. The first, Bob’s method of post-selecting the spins separately, is not available to Eve; for instance, if she post-selects in the σ_y direction, Bob will notice that the mean of σ_x changed. An alternative is then for Eve to perform a collective projection. However, in order to determine which projector she needs to measure in the 2^N -dimensional Hilbert space, Eve *must* be in possession of the code *before* she gets the spins. This information can therefore be concealed from Eve if Alice sends the code *after* the spins have passed by Eve, or if she sends the spins only one at a time. It can further be shown that by separate spin measurements, which rotate the spin by less than $\sqrt{D}$, Eve cannot gain information. The security of the protocol, hence rests on two ingredients: the non-commutativity of Bob’s final measurements, as in key distribution protocol [8], and a temporal separation between the spins and the code or between spins themselves. Such a temporal delay was suggested by Goldenberg and Vaidman [9]. However the latter proposal requires the use of a quantum storage which is not needed in our case; Bob can complete his

measurements before receiving the code.

A simple quantum optical realization of the protocols should be feasible following a previous proposal [10,11] for observing weak values through the interaction of a coherent state in a high-Q resonant cavity with a pre- and post-selected Rydberg atom. Here the photon occupation number plays the role of the pointer variable. Another realization may employ EPR-type correlations in a photon-pair generated via parametric down conversion process. In this case, one will perform a weak-measurement on one photon and an ordinary measurement on the other. Weak measurements of polarization have been suggested [12,13] and carried out for classical light [14,15]. In the present case, one step further is required, since the measurement must be performed on each photon separately.

To conclude with, it is conceptually instructive to compare the present suggestion to the Bennett-Wiesner communication scheme [16]: Alice prepares an EPR pair and sends one of the particles to Bob. When Bob wishes to send a 2-bit message to Alice, he performs a π rotation of his spin around the x , y or z axis, or does nothing. He then sends the spin back to Alice, who can reveal his actions by measuring a joint observable. This scheme, as well as other well known schemes such as teleportation [17] or quantum key distribution [8,18], ordinarily rely on entanglement and its preservation [19]. While in the present proposal the weak measurement can be formulated in terms of entanglement between the measuring device and the system, as in eq. (5), the order of events is here such that the entanglement is no longer “there” once the reading of the weak measurement has been recorded.

How should we then understand the flow of information in the present case? Bennett and Wiesner suggested after Schumacher, that since in their scheme only one qbit is returned to Alice while two bits of information are transmitted, “*one bit of information is sent forward in time... while the other bit is sent backwards in time to the EPR source, then forward in time through the untreated particle*”. In our first example, because the code is prepared before Bob sends his message, and because no useful information can be extracted from the key, the message is in some sense already “in” the code. It seems therefore that the full one bit of information is sent backwards in time. Yet, it is only with the aid of the post-selected “key” that the message is extracted from the quantum noise. Therefore, no conflict arises between macroscopic causality and this apparently retrocausal flow of information.

We thank Y. Aharonov, M. Byrd, A. Casher, T. Efron, M. Mims, Y. Ne’eman, S. Nussinov, and L. Vaidman for helpful discussions, and the Physics Department at the University of South Carolina for their kind hospitality throughout the preparation of the article. A. B. acknowledges the support of Fondo Colciencias-BID and World Laboratory. B. R. acknowledges the support from grant 471/98 of the Israel Science Foundation, established by the Israel Academy of Sciences and Humanities.

* Also at *Centro Internacional de Física, Edificio Manuel Ancizar, Universidad Nacional, Bogotá, Colombia.*

-
- [1] Y. Aharonov, D. Albert, A. Casher, and L. Vaidman, Phys. Lett. A **124** 199 (1987).
 - [2] Y. Aharonov, D. Albert, and L. Vaidman, Phys. Rev. Lett. **60** 1351 (1988).
 - [3] Y. Aharonov and L. Vaidman, Phys. Rev. A, **41**, 11 (1990).
 - [4] B. Schumacher, Phys. Rev. A **51**, 2738 (1995).
 - [5] A remarkable property exhibited by this sum rule is that the pair of weak values $\{A_w(b), A_w(b-)\}$ will generally vary with the chosen final direction $\hat{b}$, even when the probabilities $\{P(b), P(b-)\}$ remain unchanged under this variation.
 - [6] An extension of the first protocol involves distinguishing m different directions on the equal probability plane [5]. For large m the optimal number scales as m^4 , meaning that the batch size grows exponentially with the number of “directional” bits $\log_2 m$ per batch.
 - [7] This would hold, in fact, even if Alice had adopted a deception scheme in which the spins were prepared at random in either the $|x+\rangle$ or $|x-\rangle$ directions. Alice would decode the message by binning into four instead of two categories. However, Eve would still be able to decode the message from the weak averages. This is essentially a time-reversed version of a situation in which the initial states are distinguished by their corresponding expectation values.
 - [8] C. H. Bennett and G. Brassard, in *Proceedings of the IEEE International Conference on Computers, Systems, and Signal Processing, Bangalore, India* (IEEE, New York, 1984), p. 175.
 - [9] L. Goldenberg and L. Vaidman, Phys. Rev. Lett. **75**, 1239, (1995).
 - [10] Y. Aharonov, L. Davidovich, and N. Zagury, Phys. Rev. A **48** 1687 (1993)
 - [11] N. Zagury and A. F. R. de Toledo Piza, Phys. Rev. A, **50**, 2908 (1994).
 - [12] M. Duck, P.M. Stevenson, and E.C.G. Sudarshan, Phys. Rev. **D40**, 2112, (1989).
 - [13] J.M. Knight and L. Vaidman, Phys. Lett. **A143**, 357 (1990).
 - [14] N.W.M. Ritchie, J.G. Story, and R.G. Hulet, Phys. Rev. Lett. **66**, 1107 (1991).
 - [15] J. G. Story, N.W. M. Ritchie and R. G. Hulet, Mod. Phys. Lett. **B5** 1713 (1991).
 - [16] C. H. Bennett and S. J. Wiesner, Phys. Rev. Lett. **69**, 2881 (1992).
 - [17] C.H. Bennett, G. Brassard, C. Crépeau, R. Josza, A. Peres, and W.K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
 - [18] A.K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
 - [19] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. Smolin and W. K. Wootters, Phys. Rev. Lett. **76**, 722

(1996). C. Bennett, D. DiVincenzo, J.A. Smolin, and W. K. Wootters, Phys. Rev. A. **54**, 3824 (1996).